

Figure 1

PROVIDE SCAFFOLD PROTEIN BACKBONE STRUCTURE

ESTABLISH POSITION IN BACKBONE FOR HIGH ENERGY STATE ROTAMERS

ANALYZE INTERACTION OF HIGH ENERGY STATE ROTAMERS WITH PROTEIN
SCAFFOLD TO GENERATE PRIMARY LIBRARY OF VARIANT SEQUENCES WITH
PUTATIVE ENZYME-LIKE ACTIVITY

bioRxiv preprint doi: <https://doi.org/10.1101/000000>; this version posted January 1, 2014. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY-NC-ND 4.0 International license.

Figure
2






Figure
3




bioRxiv preprint doi: <https://doi.org/10.1101/000000>; this version posted January 1, 2014. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY-NC-ND 4.0 International license.

Figure
4


A.



B.




C.



20250606/004679

Figure
5



$$v = \frac{k_{cat} [\text{E}][\text{S}]}{K_m + [\text{S}]}$$

$$k_{cat} = \frac{k_2 k_3}{k_2 + k_3}$$

$$K_m = \frac{K_s k_3}{k_2 + k_3}$$

13074679 034303

Figure
6




Figure
7




Figure
8




Figure
9

